

COMPARING ABSOLUTE AND RELATIVE GROMOV–WITTEN INVARIANTS

DUSA MCDUFF

ABSTRACT. This note compares the usual (absolute) Gromov–Witten invariants of a symplectic manifold with the invariants that count the curves relative to a (symplectic) divisor D . We give explicit examples where these invariants differ even though it seems at first that they should agree, for example when counting genus zero curves in a class β such that $\beta \cdot D = 0$. The main tool is the decomposition formula in the form developed by A. Li–Ruan.

CONTENTS

1. Introduction	1
2. Relative Gromov–Witten invariants	3
2.1. Relative stable maps	3
2.2. First results on Gromov–Witten invariants	11
3. The decomposition formula	14
3.1. A genus zero example	18
References	22

1. INTRODUCTION

This is a largely expository paper about the An Li–Ruan version of relative Gromov–Witten invariants; cf. [11]. These have been used for a long time to help solve enumerative problems (cf. Ionel–Parker [8] for example) but have only recently been much used in other areas of symplectic geometry. For example, they are an essential ingredient in Hu–Li–Ruan’s work in [6] on the symplectic birational invariance of the uniruled property as well as in later papers [7, 9, 13, 18] on related topics. They are still rather little understood by symplectic geometers, even in the genus zero case.

§1 describes the moduli spaces of stable maps that enter into their definition. These involve curves on several levels as in many SFT moduli spaces. (Of course, relative Gromov–Witten invariants can be considered as a special case of SFT; cf. Bourgeois

Date: August 4, 2008.

2000 Mathematics Subject Classification. 53D45, 14N35.

Key words and phrases. relative Gromov–Witten invariants, Gromov–Witten invariants, decomposition formula, symplectic sum.

partially supported by NSF grant DMS 0604769.

et al. [1, Example 2.2].) We do not attempt to do any analysis, but rather to explain the structure of these moduli spaces through examples. §2 and §3 describe the decomposition formula and give several examples of its use.

Here are our main results.

1. We show in Remark 2.6 that when the normal bundle of the divisor D is suitably positive one can calculate genus zero Gromov–Witten invariants without involving higher level curves. (This result is greatly strengthened by Zinger in [23].) However, in genus one this need not be the case; cf. Example 2.5.

2. Our second set of results concern the contributions to the relative invariants from curves in or near D . More formally, these are the relative invariants of the pair (Y, D^+) , where $Y := \mathbb{P}(L \oplus \mathbb{C})$ is the projectivization of a complex line bundle $L \rightarrow D$ and $D^+ := \mathbb{P}(L \oplus \{0\})$ is the copy of D “at infinity” with normal bundle L^* . We denote by $D' := \mathbb{P}(\{0\} \oplus \mathbb{C})$ the copy of D along the zero section (with normal bundle L).

Understanding the relative invariants of (Y, D^+) is a crucial ingredient of the proofs of many results, such as the blow up formulas in [6, 9]. As a first step, one needs to know that certain invariants vanish. Lemma 2.8 gives a version of this result for relative invariants in class $\beta \in H_2(Y)$ under the condition that $\beta \cdot D' \geq 0$ and the homological constraints are pulled back from D . The argument is elementary, based on a dimension count. This argument fails when $\beta \cdot D' < 0$, but there were no explicit examples showing that the result may also not hold in this case. Example 2.9 gives such an example in the genus zero case.

3. Our first application of the decomposition formula is to give conditions under which the absolute and relative genus zero invariants agree for classes β such that $\beta \cdot D = 0$: see Proposition 3.2. Its statement is slightly stronger than other similar results in the literature (cf. [7, 9]) because we do not make the usual positivity assumptions on the normal bundle of D . However the argument is still based on a dimension count.

4. Finally we give examples where the absolute and relative invariants differ for classes β such that $\beta \cdot D = 0$. It is easy to find such examples in genus > 0 ; see Example 3.4. The genus zero example is trickier and is given in Proposition 3.7.

The reason why such examples exist is that in calculating the relative invariants one uses almost complex structures J and perturbing 1-forms ν that respect the divisor D . If D is “nonnegative” then a generic pair (J, ν) of this form should satisfy the regularity conditions for stable maps to X as well as those for maps to (X, D) . (This statement is made precise in Zinger [23].) However, if D is sufficiently “negative” then this is not so, and there may be contributions to the relative invariant that are perturbed away in the absolute invariant.

Acknowledgements I wish to thank the many people who have helped me understand relative invariants, in particular T.-J. Li, Yongbin Ruan, Rahul Pandharipande, and Aleksey Zinger.

2. RELATIVE GROMOV–WITTEN INVARIANTS

We first explain the structure of the stable maps used to calculate the relative invariants, illustrating by many examples. Then we describe some elementary vanishing results.

2.1. Relative stable maps. Consider a pair (X, D) where D is a divisor in X , i.e. a codimension 2 symplectic submanifold. The relative invariants count connected J -holomorphic curves in some k -fold prolongation X_k of X . Here J is an ω -tame almost complex structure on X satisfying certain normalization conditions along D . In particular D is J -holomorphic, i.e. $J(TD) \subset TD$. The invariants are defined by first constructing a compact moduli space $\overline{\mathcal{M}} := \overline{\mathcal{M}}_{\beta, \underline{d}}^{X, D}(J)$ of genus g J -holomorphic curves C in class $\beta \in H_2(X)$ as described below and then integrating the given constraints over the corresponding virtual cycle $\overline{\mathcal{M}}^{[vir]}$. Here $\underline{d} = (d_1, \dots, d_r)$ is a partition of $d := \beta \cdot D \geq 0$.

To a first approximation the moduli space consists of curves, i.e. equivalence classes $C = [\Sigma, u, \dots]$ of stable maps, that intersect the divisor D at r points with multiplicities d_i . More precisely, each such curve has $k + 1$ levels C_i for some $k \geq 0$, the principal level C_0 in $X \setminus D$ and the higher levels C_i (sometimes called bubbles) in the $\mathbb{C}^*$ -bundle $L_D^* \setminus D$ where $L_D^* \rightarrow D$ is the dual of the normal bundle to D . We identify $L_D^* \setminus D$ with the complement of the sections D_0, D_∞ of the ruled surface

$$\pi_Q : Q := \mathbb{P}(\mathbb{C} \oplus L_D^*) \rightarrow D,$$

where the zero section $D_0 := \mathbb{P}(\mathbb{C} \oplus \{0\})$ of L_D^* has normal bundle L_D^* and the infinity section $D_\infty := \mathbb{P}(\{0\} \oplus L_D^*)$ has normal bundle L_D . We then think of the original divisor $D \subset X$ as the infinity section $D_{0\infty}$ at level 0. For each $i > 0$ the ends of C along the zero section D_{i0} of the components at level i match with those along $D_{i-1\infty}$ of the preceding level, and the relative constraints are put along the infinity section $D_{k\infty}$ of the last copy of L_D^* .

Each C_i has three disjoint (and possibly empty) sets of marked points, the absolute marked points, the points $\{y_{j0} : 1 \leq j \leq r_{i0}\}$ where it meets D_{i0} and the points $\{y_{j\infty} : 1 \leq j \leq r_{i\infty}\}$ where it meets $D_{i\infty}$. (For this to make sense when $i = 0$ take D_{00} to be empty.) We require further that each component C_i be stable, i.e. have a finite group of automorphisms. For C_0 this has the usual meaning. However, when $i \geq 1$ we identify two level i curves C_i, C'_i if they lie in the same orbit of the fiberwise $\mathbb{C}^*$ action; i.e. given representing maps $u_i : (\Sigma_i, j) \rightarrow Q$ and $u'_i : (\Sigma'_i, j') \rightarrow Q$, the curves are identified if there is $c \in \mathbb{C}^*$ and a holomorphic map $h : (\Sigma_i, j) \rightarrow (\Sigma'_i, j')$ (preserving all marked points) such that $u_i \circ h = c u'_i$.¹ Thus $L_D^* \setminus D$ should be thought of as a “rubber” space; cf. [14].

Note that, although the whole curve is connected, the individual levels need not be but should fit together to form a genus g curve. The homology class β of such a curve is defined to be the sum of the homology class of its principal component with the

¹ Hence a level C_i cannot consist only of multiply covered curves $z \mapsto z^k$ in the fibers $\mathbb{C}^*$ of $L_D^* \setminus D$ because these are not stable.

projections to D of the classes of its higher levels.² When doing the analysis it is best to think that the domains and targets of the curves have cylindrical ends. However, their indices are the same as those of the corresponding compactified curves; see [11, Prop 5.3]. Thus the (complex) dimension of the moduli space $\overline{\mathcal{M}}_{g,\beta,k,(d_1,\dots,d_r)}^{X,D}$ of genus g curves in class β with k absolute marked points and r relative ones is

$$(2.1) \quad n(1-g) + c_1^X(\beta) + k + r + 3(g-1) - \sum_{i=1}^r (d_i - 1)$$

Here we subtract $d_i - 1$ at each relative intersection point of multiplicity d_i since, as far as a dimensional count is concerned, what is happened at such a point is that d_i of the $d := \beta \cdot D$ intersection points of the β -curve with D coincide.

Example 2.1. (i) Let $X = \mathbb{P}^2 \# \overline{\mathbb{P}^2}$, the one point blow up of $\mathbb{P}^2$ and set $D := E$, the exceptional curve. Denote by $\pi : X \rightarrow \mathbb{P}^1$ the projection, and fix another section H of π that is disjoint from E . Let J be the usual complex structure and $\overline{\mathcal{M}}_\lambda^{X,E}(J;p)$ be the moduli space of holomorphic lines through some point $p \in H$, where $\lambda = [H]$ is the class of a line. Since $\lambda \cdot E = 0$ there are no relative constraints. Then $\overline{\mathcal{M}}_\lambda^{X,E}(J;p)$ has complex dimension 1 and should be diffeomorphic to $\mathbb{P}^1$. The closure of the ordinary moduli space of lines in X though p contains all such lines together with one reducible curve consisting of the union of the exceptional divisor E with the fiber $\pi^{-1}(\pi(p))$ through p . But the elements of $\overline{\mathcal{M}}_\lambda^{X,E}(J;p)$ do not contain components in E . Instead, this component becomes a higher level curve lying in $Q = \mathbb{P}(\mathbb{C} \oplus \mathcal{O}(1))$ that intersects $E_0 = \mathbb{P}(\mathbb{C} \oplus \{0\})$ in the point $E_0 \cap \pi^{-1}(\pi(p))$ and lies in the class λ_Q of the line in $Q \cong X$. Note that modulo the action of $\mathbb{C}^*$ on Q there is a *unique* such bubble. Thus the corresponding two-level curve in $\overline{\mathcal{M}}_\lambda^{X,E}(J;p)$ is a rigid object. Moreover, because λ_Q projects to the class $\varepsilon \in H_2(E)$ of the exceptional divisor, its homology class $(\lambda - \varepsilon) + pr(\lambda_Q)$ is $(\lambda - \varepsilon) + \varepsilon = \lambda$. Such a bubble can also be thought of as a holomorphic section of the bundle $\mathcal{O}(1)$ that is zero at $\pi(p)$.

(ii) If X is as above but D is the line H then $\overline{\mathcal{M}}_{\lambda,1}^{X,H}(J)$ has complex dimension 2. It contains a 1-dimensional family of curves that each consist entirely of a level 1 bubble, with one such curve for each point in $H_{1\infty}$.

To state the index formula we need more notation. For simplicity let us suppose there are no absolute marked points. Then each level C_i (which may not be connected) is an equivalence class of stable maps

$$[\Sigma_i, u_i, y_{10}, \dots, y_{r_{i0}0}, y_{1\infty}, \dots, y_{r_{i\infty}\infty}],$$

in some class β_i , where the internal relative marked points $y_{10}, \dots, y_{r_{i0}0}$ are mapped to D_{i0} with multiplicities $\underline{m}_{i0}$ that sum to $\beta_i \cdot D_{i0}$ and $y_{1\infty}, \dots, y_{r_{i\infty}\infty}$ are taken to $D_{i\infty}$ with multiplicities $\underline{m}_{i\infty}$ that sum to $\beta_i \cdot D_{i\infty}$. The index I_i of such a curve is defined to

² This is the class of the curve in X obtained by gluing all the levels together. When $k = 0$ the curve only has one level and so the relative constraints lie along $D \subset X$. These are the curves counted by Ionel–Parker [8].

be the formal dimension of the moduli space of all such curves with domain of the given genus, where we impose the conditions that $u_i(y_{\ell 0}) \in D_{i0}, u_i(y_{\ell \infty}) \in D_{i\infty}$, subtract -1 (because of the $\mathbb{C}^*$ action) when $i > 0$, and also subtract $(\beta_i \cdot D_{i0} - r_{i0}) + (\beta_i \cdot D_{i\infty} - r_{i\infty})$ to take care of the multiplicity of the intersections along the divisors. The components C_i and C_{i+1} match along $D_{i\infty} = D_{i+1,0}$ only if the multiplicities $\underline{m}_{i\infty}$ and $\underline{m}_{i+1,0}$ agree (after possible reordering). Also we need corresponding ends of C_i and C_{i+1} to intersect transversally in $D_{i\infty} = D_{i+1,0}$ (so that they can be glued). Thus the index (over $\mathbb{C}$) of the $(k+1)$ -level curve $C = \cup_{i=0}^k C_i$ is

$$(2.2) \quad \text{ind } C = \sum_{i=0}^k I_i - r(n-1),$$

where I_i is as above and $r = \sum_{i=1}^k r_i$ is the number of internal intersection points.

The next result was noted by Ionel–Parker [8, Prop. 14.10].

Lemma 2.2. *In the genus zero case, the index of a higher level component in Q equals that of its projection to D (considered as a curve in D).*

Proof. Consider a sphere in Q in a class $\beta := \alpha + df$, where f is the class of the fiber of $\pi_Q : Q \rightarrow D$ and $\alpha \in H_2(D_0)$, that intersects D_∞ at r_∞ points through cycles $b_\infty := (b_{\infty i})$ with multiplicities $d_{\infty i}$ summing to d , and intersects D_0 at r_0 points through cycles $b_0 := (b_{0j})$ with multiplicities d_{0j} summing to $d + \alpha \cdot D_0$. Let δ_b be half the sum of the degrees of the cycles $b_{\infty i}, b_{0j}$ and set $r := r_\infty + r_0$, the total number of constraints. Then the formal (complex) dimension of the moduli space of such spheres (when quotiented out by the $\mathbb{C}^*$ action on Q) is

$$\begin{aligned} n + (c_1^D(\alpha) + \alpha \cdot D_0 + 2d) + (r-3) - (d-r_\infty) - (d + \alpha \cdot D_0 - r_0) + (\delta_b - rn) - 1 \\ = n + c_1^D(\alpha) + 2r - 3 + \delta_b - rn - 1. \end{aligned}$$

Here $n := \dim_{\mathbb{C}} X$, $c_1^D(\alpha) + \alpha \cdot D_0 + 2d = c_1^Q(\beta)$, the term $r-3$ takes care of the variation of the marked points, the terms $d-r_\infty, d + \alpha \cdot D_0 - r_0$ appear because we must subtract $d_i - 1$ at each relative insertion of order $d_i > 1$, and we subtract $rn - \delta_b$ to take care of the homological constraints.

When we project these spheres to D they still have r marked points through the constraints $b_{\infty i}, b_{0j}$, now considered as lying in D , but all information about the orders of the normal tangencies is lost. The formal dimension of the moduli space of such curves is $(n-1) + c_1^D(\alpha) + r - 3 + \delta_b - r(n-1)$, just what we found above. $\square$

The next result can be easily checked; it is implicit in Li–Ruan [11] formula (5.1).

Lemma 2.3. *The index of a genus g curve in class β with $k+1$ levels and given incidence multiplicities along $D_{\infty k}$ is $I(g, \beta) - k$, where $I(g, \beta)$ is the index of the main stratum of curves with no higher levels.*

The constraints for the relative invariants consist of homology classes b_j in the divisor D (the relative insertions) together with absolute (descendent) insertions $\tau_{i_j} a_j$ where

$a_j \in H_*(X)$ and $i_j \geq 0$. We shall denote the (connected) relative genus zero invariants by:

$$(2.3) \quad \langle \tau_{i_1} a_1, \dots, \tau_{i_q} a_q \mid b_1, \dots, b_r \rangle_{0, \beta, (d_1, \dots, d_r)}^{X, D},$$

where $a_i \in H_*(X)$, $b_i \in H_*(D)$, and $d := \sum d_i = \beta \cdot D \geq 0$. (If $\beta \cdot D < 0$ then the moduli space is undefined and the corresponding invariants are set equal to zero.) This invariant counts connected genus zero curves in class β that go through cycles representing the classes a_i and intersect D to order $\underline{d} := (d_1, \dots, d_r)$ in the b_i , i.e the i th relative marked point intersects D to order $d_i \geq 1$ at a point on some representing cycle for b_i . We assume that the cycles a_i are in general position to D , while the b_i lie in D . Moreover, the insertion τ_i occurring at the j th absolute marked point means that we add the constraint $(c_j)^i$, where c_j is the first Chern class of the cotangent bundle to the domain at the j th marked point. One can evaluate (2.3) (which in general is a rational number) by integrating an appropriate product of Chern classes over the virtual cycle corresponding to the moduli space of stable J -holomorphic maps that satisfy the given homological constraints and tangency conditions. For details on how to construct this virtual cycle see for example Li–Ruan [11] or Hu–Li–Ruan [6].³ Since the virtual cycle has dimension equal to the relevant index, the above formula defines a number only when this index equals the sum of the dimensions of the constraints. In all other cases the invariant is defined to be zero.

We now give some examples to explain the structure of the stable maps considered here, and to illustrate the index calculations.

Example 2.4. (i) Let $(X, D) = (\mathbb{P}^2, \mathbb{P}^1)$ and consider the moduli space $\overline{\mathcal{M}}_{2\lambda, (2), 4}^{X, D}(p) =: \overline{\mathcal{M}}$ of degree 2 stable maps, with 3 absolute marked points z_1, z_2, z_3 and one relative marked point z_0 , and that are tangent to D at the fixed point $p = u(z_0) \in D$. Its open stratum $\mathcal{M}$ consists of equivalence classes $[u; z_0, \dots, z_3]$ where $u : \mathbb{P}^1 \rightarrow \mathbb{P}^2$ is a map of degree 2 (i.e. a conic) such that $u^{-1}(p) = u^{-1}(D) = z_0$. The (complex) dimension of $\mathcal{M}$ is 6, one less than the dimension of the space of such maps with $u(z_0) = p$, since the tangency condition can be interpreted as saying that the two intersection points of the conic with D coincide. The evaluation map takes this open stratum injectively onto the set of generic triples of points $u(z_1), u(z_2), u(z_3)$ in $\mathbb{P}^2$.

The full moduli space $\overline{\mathcal{M}}$ contains various degenerate curves. We now investigate some of the corresponding strata, showing that their dimensions agree with the formula in Lemma 2.3. The usual stratification of $\overline{\mathcal{M}}$ (which is determined by the domain and marked points of the stable map) is closely related to the positions of the image points $u(z_1), u(z_2), u(z_3)$, and we shall analyze $\overline{\mathcal{M}}$ by looking at this image configuration rather than the strata.

³ What one needs here is an appropriate framework in which to construct suitable multivalued perturbations as in [17]; see also [16]. There are many possible ways of solving this problem. A very general construction that applies in a wide variety of situations will eventually be provided by Hofer–Wysocki–Zehnder [4] and [3]. This should give a coherent setting in which one could reprove the results claimed here. For another approach see Zinger [22, 23].

If $u(z_1), u(z_2), u(z_3)$ are collinear the curve splits into a line ℓ in $\mathbb{P}^2$ through these three points together with a bubble component in $Q := \mathbb{P}(\mathbb{C} \oplus L_D^*)$ through the point $q := \ell \cap D$ on the zero section $D_0 := \mathbb{C} \oplus \{0\}$ of Q and tangent to the infinity section D_∞ at the point $u(z_0)$. One can think of this bubble as a meromorphic section of L_D^* with a pole of order 2 at p and a simple zero at q , or from a more symplectic viewpoint as a curve in the compact ruled surface Q in the class $D_0 + 2f$, where f denotes the class of the fiber of $Q \rightarrow D$. It is counted modulo the $\mathbb{C}^*$ action that fixes 0 and the points of D_∞ . Since Q is the one point blow up of $\mathbb{P}^2$ with exceptional divisor D_0 , each such bubble corresponds to a conic on $\mathbb{P}^2$ (the blow down of Q) that is tangent to D_∞ at a fixed point on D_∞ and also tangent to a fixed direction at 0. There is one such conic modulo the $\mathbb{C}^*$ action. Hence this bubble is rigid (i.e. lies in a zero-dimensional moduli space) and contributes to the nonzero rubber invariant

$$\langle pt | pt \rangle_{0, 2f + \lambda_0, (1), (2)}^{D_0, Q, D_\infty, \sim}$$

that counts curves with these constraints modulo the $\mathbb{C}^*$ action on the fibers of $Q \rightarrow D$. (Here the relative constraints are put on the left and right of the bracket and we use $\sim$ to denote rubber invariants as in Maulik–Pandharipande [14].) Since the (collinear) points $u(z_1), u(z_2), u(z_3)$ have 5 degrees of freedom, these curves form a codimension 1 subspace of the full moduli space. This is consistent with Lemma 2.3; these curves are regular and so should form a space whose dimension equals the index of the given stratum.

A second kind of degeneration occurs when $u(z_1)$ and $u(z_2)$ lie on a line through the fixed point p . Now the principal part of the curve has two components, one a line through $u(z_1), u(z_2)$ and the other a line through $u(z_3)$ and p , that are joined by a level 1 bubble in class $2f$ that lies in the fiber over p , intersecting D_{10} twice at p with order 1 and intersecting $D_{1\infty}$ once with order 2; see Figure 2.1. This fiberwise bubble is rigid, and contributes to the nonzero rubber invariant

$$\langle D_0, D_0 | pt \rangle_{0, 2f, (1,1), (2)}^{D_0, Q, D_\infty, \sim}.$$

Because this rubber invariant counts curves in a fiber class that go through a fixed point, it equals the corresponding invariant in the fiber of $Q \rightarrow D$. Identify this fiber with $S^2 := \mathbb{C} \cup \{\infty\}$, meeting D_0, D_∞ in the points $\{0\}$ and ∞ . Then this invariant counts the maps $\mathbb{C} \rightarrow \mathbb{C}$ of the form $z \mapsto az(z - b)$ modulo the reparametrization $z \mapsto cz$ of the domain and the $\mathbb{C}^*$ action $w \mapsto dw$ on the target. Hence this invariant is 1. Again the points $u(z_1), u(z_2), u(z_3)$ have 5 degrees of freedom, so that this stratum of regular curves has codimension 1.

If all 4 points are collinear (but still distinct), then we could have:

- (a) two distinct lines through p together with a bubble at p as in the preceding paragraph but with z_1, z_2, z_3 all on the same line,
- (b) two coincident lines plus a bubble (with various possibilities for the marked points),
- or
- (c) a doubly covered line through these 4 points with one of its branch points at p .




FIGURE 2.1. The two component stable map in Example 2.4 (i) when $p, u(z_1)$ and $u(z_2)$ are collinear. The schematic picture of the bubble is meant to illustrate its parametrization; its image lies entirely in the fiber over p and it intersects the divisors $D_{10}, D_{1\infty}$ with multiplicities $(1, 1), (2)$.

Since the collinearity condition has index 2 we would expect the corresponding spaces of stable maps to have codimension 2. This is true in case (b), because although the curves are regular and have two levels their principal components satisfy an extra constraint. However in cases (a) and (c) the curves have 5 degrees of freedom; e.g. in (c) there is one for the line, one for its second branch point and three for the positions of the other points on it. In these degenerate cases the curves are not uniquely determined by the position of the points $u(z_1), u(z_2), u(z_3)$, more precisely the collinearity conditions are not transverse to the evaluation map $\overline{\mathcal{M}} \rightarrow (\mathbb{P}^2)^3$ and so do not cut down its dimension in the expected way.

Other degenerations occur when some of the points coincide; we leave further discussion to the reader.

(ii) Now consider the moduli space $\overline{\mathcal{M}}$ of conics in $\mathbb{P}^2$ with 3 absolute and 2 relative marked points that intersect the line D with multiplicity $\underline{2} = (1, 1)$. This has dimension 8. $\overline{\mathcal{M}}$ contains a codimension 1 stratum $\mathcal{S}$ consisting of curves whose two intersection points with D coincide. The open stratum in $\mathcal{S}$ is the 7-dimensional space of two-level curves where C_0 is tangent to $D_{0\infty}$ at some unspecified point and C_1 is a rigid fiberwise bubble contributing to the rubber invariant

$$\langle pt | D_{\infty}, D_{\infty} \rangle_{0, 2f, (2), (1, 1)}^{D_0, Q, D_{\infty}, \sim}.$$

In other words, C_1 is tangent to D_{10} at some specified point and intersects $D_{1\infty}$ twice transversally at arbitrary points. Thus the incidence conditions of the bubble along $D_{1\infty}$ have type $(1, 1)$ while the incidence conditions along the internal divisor D_{10} are dual to those of the principal component. Note that these are not the same as the elements considered in (i) because the incidence conditions along D have type $(1, 1)$ instead of (2) .




FIGURE 2.2. Some of the stable maps considered in Example 2.5. In (I) S is an immersed sphere of degree 3 with its node on $D_{0\infty}$ while in (II) it is tangent to D . Each bubble lies in a single fiber. Hence when a bubble intersects one of the divisors D_{ij} with multiplicity $(1,1)$ its two intersection points with D_{ij} coincide.

Example 2.5. Consider the moduli space of degree 3 tori in $X := \mathbb{P}^2$ that intersect $D := \mathbb{P}^1$ with multiplicity $(2,1)$. Its main stratum consists of tori tangent to D and by formula (2.1) has dimension 8. Figure 2.2 illustrates two ways in which the genus of the stable map may be carried by the pattern of intersections of the bubble components, rather than by the genus of any single component.

In (I), the main component S is an immersed sphere of degree 3 that intersects D transversally at its node as well as at one other point. The level 1 curve consists of two fiberwise bubbles, one of which is doubly covered and intersects D_{10} with multiplicity $(1,1)$, thus creating some genus. Since both bubbles are rigid, the dimension of the space of curves of this type is the same as the dimension of the space of spheres S with node on D ; thus it is 7, which agrees with the formula in Lemma 2.3.

In (II) the main component is still an immersed sphere of degree 3, but now we assume that it is tangent to D at some point x , and has one other transverse intersection point with D . In particular, its node is disjoint from D . There are two other levels that contain doubly covered spheres C_1^x, C_2^x in the fiber over x that are tangent to D_{10} and $D_{2\infty}$ but intersect the intermediate divisors $D_{1\infty}, D_{20}$ with multiplicity $(1,1)$. (Each of these levels also contains a singly covered sphere through the other intersection with D , but these are irrelevant to the present discussion.) The dimension of this stratum is again equal to the dimension of the family of spheres C_0 , and so is 7, which is not equal to the index. The point is that although the components C_i^x are stable and regular, they lie in moduli spaces that do *not* intersect transversally along $D_{1\infty} = D_{20}$ because their two intersections with this divisor must always coincide. Therefore the only way to regularize this stratum is by gluing. The dimensions then work out correctly, because one must subtract 1, the dimension of the relevant obstruction bundle. One way to understand this is to note that the two bubbles C_1^x and C_2^x do glue to give a family of regular tori in the single $\mathbb{P}^1$ fiber over $x \in D$. However this torus has trivial normal bundle when considered as a curve in Q and so is *not* regular in Q but rather has a

cokernel of dimension $\dim D$. This remark also explains why there is no similar problem with the index in case (I) since in this case the glued curve lies in X .

The above nontransversality phenomenon does not occur in genus zero. As we now explain, it follows that as long as the divisor D is suitably positive we can ignore the curves with higher levels.

Remark 2.6. Suppose that D has “nonnegative” normal bundle L_D ; for example we might suppose that $c_1(L_D) = \lambda[\omega|_D]$ for some $\lambda \geq 0$, or, more generally that $c_1(L_D)(\alpha) \geq 0$ for all classes $\alpha \in H_2(D)$ that can be represented by J_D -holomorphic spheres. Then in the genus zero case, *curves with higher levels do not contribute to relative invariants*

$$(2.4) \quad \langle a_1, \dots, a_q \mid b_1, \dots, b_r \rangle_{0, \beta, (d_1, \dots, d_r)}^{X, D}$$

with only homological constraints. By this we mean the following. Denote by $\overline{\mathcal{M}}_{\beta, \underline{d}}^{X, D}(J)$ the moduli space of all genus 0 stable J -holomorphic curves in class β with q absolute and r relative marked points. Suppose we partially regularize this moduli space by perturbing the equations for the components of the curves in each level to make them as generic as possible, but not performing any gluing operations. Here we only allow perturbations ν that preserve the structure of the target; i.e. we look at equations of the form $\bar{\partial}_J u = \nu(u)$ where the perturbing 1-form $\nu(u)$ at level 0 vanishes sufficiently fast in the normal directions to D and at higher levels is pulled back from D by the projection $\pi : Q \rightarrow D$. Hence each higher level component still projects to a curve in D and our procedure regularizes each of these projected components as a curve in D .

We claim that in the genus 0 case this procedure is enough to reduce the dimension of the set $\mathcal{S}_H$ of curves with higher levels below that of the full moduli space. Hence the partially regularized set $\mathcal{S}_H$ does not have large enough dimension to go through all the homological constraints and so does not contribute to the invariant. This is not true in higher genus because of the possible existence of degenerate objects such as the nontransverse multiply covered fibers of Example 2.5.

There are several points to note about the proof of the claim. Denote by $C_i^D, i = 1, \dots, \ell$, the projections to D of the higher level components of C . The adjustments to the index coming from the multiplicities $\underline{d}$ do not affect the argument, and so for simplicity we will ignore them, assuming in effect that $\underline{d} = (1, \dots, 1)$. Then, the relative index is just $\text{ind}_X \beta$, the index of genus zero β curves in X .

First note that, by Lemma 2.2, our perturbations are sufficient to reduce the dimension of the moduli space of each C_i^D to its index (in D). Secondly, we may arrange that the C_i^D intersect transversally in D because their intersection pattern is a tree so that two distinct components C_i^D, C_j^D have at most one intersection point; cf. [20, §6.3]. Thirdly, because of the positivity assumption $c_1(L_D)(C_i^D) \geq 0$ for all i , the curves C_i^D are regular when considered as curves in X , and (because they all can be perturbed off D) their intersections are transverse in X as well as in D .

Now suppose that $\ell = 1$ and C has no components at level zero. Then C consists of single component in Q whose projection to D lies in some class β , and the claim holds

because $c_1^D(\beta) \leq c_1^X(\beta)$, so that $\text{ind}_D \beta < \text{ind}_X \beta$. On the other hand, if C has more than one component, it follows readily from the above remarks show that its index in the relative moduli space is at most equal to the index of the stable map in X with components C_0, C_i^D and so is less than $\text{ind}_X \beta$. Hence if C has higher levels it does not lie in the top dimensional stratum and so cannot satisfy all the homological constraints in (2.4).

The failure of the above argument for “negative” D is the reason why absolute and relative GW invariants can differ in cases when one might think they should agree, e.g. if $\beta \cdot D = 0$; cf. Proposition 3.7.

2.2. First results on Gromov–Witten invariants. We now explain some vanishing results that follow by dimensional arguments. As a warmup, we begin by considering the relative invariants of $(\mathbb{P}^n, \mathbb{P}^{n-1})$.

Lemma 2.7. *Let $X = \mathbb{P}^n$ and $D = \mathbb{P}^{n-1}$ be the hyperplane. Denote by λ the class of a line. Suppose that $n > 1$ and $d > 0$ or $n = 1$ and $d > 1$. Then*

$$\langle |b_1, \dots, b_r\rangle_{0, d\lambda, \underline{d}}^{\mathbb{P}^n, \mathbb{P}^{n-1}} = 0, \quad \text{for any } b_i \in H_*(\mathbb{P}^{n-1}).$$

Proof. We show that under the given conditions on n and d it is impossible to choose $\underline{d}$ and the b_i so that $\overline{\mathcal{M}}_{d\lambda, \underline{d}}^{\mathbb{P}^n, \mathbb{P}^{n-1}}$ has formal dimension 0. Therefore the invariant vanishes by definition.

By equation (2.1) the formal (complex) dimension of the moduli space of genus 0 stable maps through the relative constraints $b_1, \dots, b_r$ is

$$n + d(n+1) + r - 3 - (d-r) + \delta - rn,$$

where δ is half the sum of the degrees of the b_i . Thus $0 \leq \delta \leq rn$. Since $d-r \geq 0$ and $r > 0$, we therefore need $(d-r)n + n + 2r + \delta = 3$, which implies $d = r$ and $n = r = 1$. This contradicts our assumptions on n and d . $\square$

Our next result concerns manifolds that are fibrations with fiber $\mathbb{P}^1$. These are important because, as we shall see in §3 they can be used to understand the relation between the absolute and relative invariants. There are much more complete discussions of this topic in Maulik–Pandharipande [14], Hu–Ruan [7] and T.-J. Li–Ruan [13].

Let $L \rightarrow D$ be a complex line bundle and $Y := \mathbb{P}(L \oplus \mathbb{C})$ be its projectivization. It contains two divisors that we call D', D^+ , where $D' := \mathbb{P}(\{0\} \oplus \mathbb{C})$ is the copy of D along the zero section of L and $D^+ := \mathbb{P}(L \oplus \{0\})$ is the copy “at infinity”. Thus D' has normal bundle L while D^+ has normal bundle L^* .⁴

Given $\alpha \in H_2(D)$, we shall write α' for the corresponding element of $H_2(D')$, considered where appropriate as an element in $H_2(Y)$. Let $f \in H_2(Y)$ be the fiber class of the projection $\pi : Y \rightarrow D$.

Lemma 2.8. *Let $\beta \in H_2(Y)$ and $r, s \geq 0$.*

⁴ There are many commonly used notations; we are using that of [18] where the formula are applied to a divisor D that is a blow up so that its normal bundle L is considered as “negative”.

(i) If $\beta \cdot D' \geq 0$ but $\beta \neq \ell f$, then

$$\langle a_1, \dots, a_s | b_1, \dots, b_r \rangle_{0, \beta, \underline{d}}^{Y, D^+} = 0,$$

for any $b_i \in H_*(D^+)$ and any $a_j \in H_*(Y)$ that can be represented by cycles of the form $\pi^{-1}(a'_j)$ where $a'_j \subset D$.

(ii) If $\beta = \ell f$ then the invariant is nonzero only if $\ell = 1$, and $s = 0$.

Proof. Write $\beta = \ell f + \alpha'$ where $\ell \geq 0$. Suppose first that there are no absolute constraints and that $\alpha' \neq 0$. Choose an ω -tame almost complex structure J on Y that is invariant under the S^1 action in the fibers of the ruled manifold $Y \rightarrow D$. The fixed point set of S^1 consists of the divisors D' , D^+ , and hence it fixes the relative constraints. The localization principle⁵ implies that the only contributions to $\langle |b_1, \dots, b_r \rangle_{0, \beta, \underline{d}}^{Y, D^+}$ come from S^1 -invariant elements of the moduli space of J -holomorphic curves. Such curves would have to consist of r (possibly multiply covered) spheres in the fiber of $Y \rightarrow D$ together with a connected genus zero curve in D' . Since $\beta \cdot D' = \ell + \alpha' \cdot D' \geq 0$, we must have $\alpha' \cdot D' \geq -\ell$. Hence $c_1^Y(\beta) \geq \ell + c_1^{D'}(\alpha) =: \ell + c'$ and so, because we always count curves of dimension zero, equation (2.1) implies that

$$n + \ell + c' + r - 3 + \delta - (\ell - r) \leq rn,$$

where δ is the sum of half the dimensions of the constraints b_i .

Now observe that we may regularize the moduli space $\overline{\mathcal{M}}_{\alpha'}^{D'}$ of α' -curves (considered as curves in D') by perturbations that are tangent to D' and hence are S^1 invariant. For the given relative invariant to be nonzero the space of S^1 -invariant stable maps made up of such partially regularized α' -curves plus some fibers going through the constraints must be nonempty. Hence there must be generic α' -spheres in D' with r marked points that meet the fibers through $b_1, \dots, b_r$. In other words, we need $n - 1 + c' + r - 3 \geq r(n - 1) - \delta$. But this contradicts the inequality found above.

Now suppose that there are some absolute constraints. Since these may be represented by cycles that are invariant under the S^1 action, S^1 still acts on the space of all J -holomorphic stable maps through the constraints. Hence the localization principle again says that the invariant is zero unless this moduli space has fixed points. These would consist of some fibers through the relative constraints as well as some of the absolute constraints, plus some α' -curves in D' meeting all these fibers and going through the other absolute constraints. Hence the moduli space of partially regularized α' -curves in D' must meet all the b_i as well as the classes a'_j . A dimension count similar to that above shows that this is impossible. This proves (i).

In case (ii) we are counting ℓ -fold coverings of the fiber 2-spheres. The number of fibers through the constraints is finite and nonzero only if the intersection product in $H_*(D)$ of all the classes b_i and a'_j is nonzero and lies in $H_0(D)$. For each point in this

⁵ [19] gives a proof for symplectic nonrelative invariants. The same arguments work for relative invariants.

intersection, we are then reduced to a fiber invariant of the form

$$\langle \mathbb{P}^1, \dots, \mathbb{P}^1 | pt, \dots, pt \rangle_{0, \ell, f, \underline{d}}^{\mathbb{P}^1, pt},$$

where the absolute constraints are s copies of $\mathbb{P}^1 = a_i \cap \pi^{-1}(pt)$ and the relative ones are r points. If $s > 0$ such an invariant is nonzero only if the total number $s + r$ of constraints is ≤ 3 . For then the indeterminacy of the absolute constraint(s) can be absorbed by reparametrizations so that there is a finite number of equivalence classes of stable maps satisfying the given conditions. A dimension count then shows that $\ell = 1$. If $s = 0$ the invariant vanishes when $\ell > 1$ by Lemma 2.7. $\square$

Example 2.9. The vanishing result in Lemma 2.8 need not hold when $\beta \cdot D' < 0$. To see this, let $D = \mathbb{P}^3 \# 2\bar{\mathbb{P}}^3$ be the 2-point blow up of $\mathbb{P}^3$. Denote the classes of $\mathbb{P}^2$ and the two exceptional divisors by $H, E_1, E_2 \in H_4(D)$; write $\lambda := [\mathbb{P}^1] = H \cdot H$ and $\varepsilon_i := E_i \cdot E_i$ for the corresponding generators of $H_2(D)$. (Thus ε_i is represented by a line in $E_i \cong \mathbb{P}^2$.) Let L be the line bundle over D whose first Chern class c_1^L is Poincaré dual to $H - E_1 - E_2$. (Thus L is the normal bundle of D in the two point blow up of $\mathbb{P}^4$ when D is identified with the proper transform of a hyperplane through the two blown up points.) Let $Y = \mathbb{P}(L \oplus \mathbb{C})$ as above, with the obvious complex structure J_0 .

Consider the class $2\alpha' + f \in H_2(Y)$ where $\alpha' = \lambda' - \varepsilon'_1 - \varepsilon'_2 \in H_2(D')$. We claim that for all $\rho \in H_4(D)$

$$(2.5) \quad \langle |\rho \rangle_{0, 2\alpha' + f, (1)}^{Y, D^+} = \frac{\rho \cdot_D \alpha}{4}.$$

where $\alpha \in H_2(D)$ corresponds to α' and $\cdot_D$ denotes the intersection product in D . One could prove this by using algebro-geometric localization, but we shall take another approach.

As in Lemma 2.8, it suffices to calculate the contribution to this invariant from the moduli space $\mathcal{V}$ of $\mathbb{C}^*$ -invariant J_0 -holomorphic $(2\alpha' + f)$ -curves in Y , where J_0 denotes the obvious complex structure on Y . The elements of $\mathcal{V}$ consist of doubly covered α' -spheres in D' plus a fiber. We can regularize the space of doubly covered α' -spheres in D' , obtaining a virtual moduli cycle $\overline{\mathcal{M}}_{2\alpha'}(D')^{[vir]}$ consisting of a finite number of embedded spheres $C' = \text{im } u'$ in D' that each have a rational weight and are (J_0, ν) -holomorphic, i.e. they satisfy a perturbed Cauchy–Riemann equation of the form $\bar{\partial}_{J_0} u' = \nu(J_0, u')$. The sum of these weights is well known to be

$$\langle \rangle_{0, 2\alpha}^{D'} = \langle \rangle_{0, 2(\lambda - \varepsilon_1 - \varepsilon_2)}^{\mathbb{P}^3 \# 2\bar{\mathbb{P}}^3} = 1/8,$$

see for example Gathmann [2, Ex 8.5]. Therefore, we need to see that each element $u' \in \overline{\mathcal{M}}_{2\alpha'}(D')^{[vir]}$ gives rise to two curves in Y in class $2\alpha' + f$ and through ρ . This may be done by using the gluing methods developed by Zinger [21, §3]. We shall give a more elementary argument that exploits the fact that Y is an S^2 -bundle over D .

The perturbing 1-form ν used to construct $\overline{\mathcal{M}}_{2\alpha'}(D')^{[vir]}$ is a (multivalued) section of the (orbi)bundle $\mathcal{E}(D) \rightarrow \mathcal{W}(D)$, where $\mathcal{W}(D)$ is a neighborhood of the space of J_0 -holomorphic $2\alpha'$ -spheres in the space of all stable maps into D in class $2\alpha'$, and $\mathcal{E}(D) \rightarrow \mathcal{W}(D)$ is the (orbi)bundle whose fiber at $[\Sigma, v]$ is the space of antiholomorphic

1-forms on the nodal Riemann surface Σ with values in $v^*(TD)$. There is a similar bundle $\mathcal{E}(Y) \rightarrow \mathcal{W}(Y)$ for stable maps in class $2\alpha' + f$. The projection $\pi : Y \rightarrow D$ induces a map $\mathcal{W}(Y) \rightarrow \mathcal{W}(D)$, and also for each stable map $[\Sigma, v] \in \mathcal{W}(Y)$ a surjective bundle map $\pi_* : v^*(TY) \rightarrow (\pi \circ v)^*(TD)$.

Consider the space Γ^ν of (multi)sections ν^Y of $\mathcal{E}(Y) \rightarrow \mathcal{W}(Y)$ that are ν -compatible, in the sense that

$$\pi_*(\nu^Y(v)) = \nu(\pi \circ v), \quad \text{for all } v \in \mathcal{W}(Y).$$

Then, for every (J_0, ν^Y) -holomorphic map $v : \Sigma \rightarrow Y$, the composite $\pi \circ v$ is a (J_0, ν) -holomorphic map to D . Therefore each (J_0, ν^Y) -holomorphic sphere in $\mathcal{W}(Y)$ lies over a (J_0, ν) -holomorphic sphere in $\mathcal{W}(D)$, i.e. an element in $\overline{\mathcal{M}}_{2\alpha'}(D')^{[vir]}$. This reduces the problem of counting the (J_0, ν^Y) -holomorphic spheres in Y through the constraint ρ to a problem in the 4-dimensional spaces $X_{u'} := u'^*(\pi^{-1}(C'))$, where $C' := \text{im } u' \cong S^2$. But $X_{u'} \cong S^2 \times S^2$ and the class $2\alpha' + f$ pulls back to $[pt \times S^2]$. Moreover, one can check directly that the allowed perturbations of ν^Y are sufficient to regularize the curves in Y ; more precisely, for generic ν^Y , every (J_0, ν^Y) -holomorphic sphere in $X_{u'}$ is regular both in $X_{u'}$ and also (because u' is regular) in Y . Hence the number of such spheres though a generic point in $X_{u'}$ is 1. Since $\rho \cdot_Y [\pi^{-1}(C')] = 2\rho \cdot_D \alpha$, the result follows.

3. THE DECOMPOSITION FORMULA

The main tool in the theory of relative invariants is the decomposition rule of Li–Ruan [11] and (in a slightly different version) of Ionel–Parker [8]. So suppose that the manifold (M, Ω) is the fiber sum of (X, D, ω_X) with (Y, D^+, ω_Y) , where the divisors (D, ω_X) and (D^+, ω_Y) are symplectomorphic with dual normal bundles. This means that there is a (real) codimension 1 hypersurface S in (M, ω) such that the orbits of its characteristic foliation form a fibration of S by circles such that the quotient space can be identified with (D, ω_X) . We will only consider cases in which S divides M into two components, one symplectomorphic to $X \setminus D$ and the other to $Y \setminus D^+$.

Let us first assume that the absolute constraints can be represented by cycles in M that do not intersect S , i.e. that $a_i \in H_*(X \setminus D)$, $i \leq q$, and $a_i \in H_*(Y \setminus D^+)$, $q < i \leq p$. For simplicity we will assume throughout that the map $H_2(M) \rightarrow H_2(X \cup_D Y)$ is injective. (This hypothesis is satisfied whenever $H_1(D) = 0$, and means that there are no rim tori (cf. [8]).) Further, let $b_i, i \in I$, be a basis for $H_*(D) = H_*(D; \mathbb{Q})$ with dual basis b_i^* for $H_*(D)$. Then the decomposition formula has the following shape:

$$(3.1) \quad \langle a_1, \dots, a_p \rangle_{g, \beta}^M = \sum_{\Gamma, \underline{d}, (i_1, \dots, i_r)} n_{\Gamma, \underline{d}} \langle a_1, \dots, a_q \mid b_{i_1}, \dots, b_{i_r} \rangle_{\beta_1, \underline{d}}^{\Gamma_1, X, D} \langle a_{q+1}, \dots, a_p \mid b_{i_1}^*, \dots, b_{i_r}^* \rangle_{\beta_2, \underline{d}}^{\Gamma_2, Y, D^+}.$$

Here we sum with rational weights $n_{\Gamma, \underline{d}}$ over all decompositions $\underline{d}$ of d , all possible connected labelled trees Γ , and all possible sets $i_1, \dots, i_r$ of relative constraints. Each Γ describes a possible combinatorial structure for a stable map that glues to give a β -curve of genus g . Thus Γ is a disjoint union $\Gamma_1 \cup \Gamma_2$, where the graph Γ_1 (resp. Γ_2)

describes the part of the curve lying in some X_k (resp. some Y_ℓ) and β_1 (resp. β_2) is the part of its label that describes the homology class. Hence the pair (β_1, β_2) runs through all decompositions such that the result of gluing the two curves in the prolongations X_{k_1} and Y_{k_2} along their intersections with the relative divisors gives a curve in class β . Moreover, there is a bijection between the labels $\{(d_i, b_i) \in \mathbb{N} \times H_*(D)\}$ of the relative constraints in Γ_1 and those $\{(d_i, b_i^*) \in \mathbb{N} \times H_*(D^+)\}$ in Γ_2 . (These labels are called relative “tails” in [6].) Γ_i need not be connected; if it is not, we define $\langle \dots | \dots \rangle^{\Gamma_i}$ to be the product of the invariants defined by its connected components.

Remark 3.1. (i) In the genus zero case, each component of Γ_1 is a tree and has at most one relative tail in common with each component of Γ_2 . In many cases one can show that one side is connected, so that the other side has r components, one for each relative constraint.

(ii) The number $n_{\gamma, \underline{d}}$ is the reciprocal of the number of automorphisms of the corresponding labelled graph. We shall not be more specific here, because it will be 1 in all cases where we shall need to know its value.

(iii) If absolute constraints $a_i \in H_*(M)$ cannot be represented on one side of S or the other, one must represent each a_i by a cycle that decomposes as $a_i^X \cup a_i^Y$, where a_i^X, a_i^Y are cycles in the closure of the appropriate component of $M \setminus S$ with equal boundaries $a_i^X \cap S = a_i^Y \cap S$ on S that are unions of S^1 fibers. Then in (3.1) one must also sum over all subsets $I \subset \{1, \dots, n\}$, where the Γ_1 -curve goes through the absolute constraints $a_i^X, i \in I$ and the Γ_2 -curve goes through $a_j^Y, j \notin I$. If there are symmetries among these constraints (eg if $a_1^X = a_2^X$), one must adjust the multiplicities n_Γ accordingly.

The decomposition formula can be used to compare absolute and relative invariants. Part (i) of the following result was proved in Hu [5] and Gathmann [2] (in the projective case) under slightly weaker assumptions. Also see Lai [9].

Proposition 3.2. *Suppose that D is a divisor in (X, ω) such that for some $\beta \in H_2(X)$*

$$c_1^X(\alpha) > (n-3)(k-2) \text{ if } n \geq 3, \quad c_1^X(\alpha) > 0 \text{ if } n = 2,$$

for all $\alpha \in H_2(D)$ with $0 < \omega(\alpha) < \omega(\beta)$ and $\alpha \cdot D = -k < -1$. Then:

(i) *If $\beta \cdot D = 0$, then for any $a_i \in H_*(X \setminus D)$,*

$$\langle a_1, \dots, a_k | \rangle_{0, \beta}^{X, D} = \langle a_1, \dots, a_k \rangle_{0, \beta}^X.$$

(ii) *If $\beta \cdot D > 0$, then for any $a_i \in H_*(X \setminus D)$*

$$\langle a_1, \dots, a_k | D, \dots, D \rangle_{0, \beta, (1, \dots, 1)}^{X, D} = \langle a_1, \dots, a_k \rangle_{0, \beta}^X.$$

Proof. (i) Let L be the normal bundle of D . Decompose X into the sum of (X, D) with the ruled manifold (Y, D^+) where $Y := \mathbb{P}(L \oplus \mathbb{C})$ and $D^+ := \mathbb{P}(L \oplus \{0\})$, so that D^+ has normal bundle L^* . We put the constraints into X . There is one term in the decomposition formula with $\Gamma_2 = \emptyset$ which contributes $\langle a_1, \dots, a_k | \rangle_{0, \beta}^{X, D}$ to the absolute invariant. Hence we need to see that there are no other terms.

Observe first that, even if $k = 0$ it is impossible for there to be a nonzero term with $\Gamma_1 = \emptyset$. For then the Γ_2 -invariant must be $\langle | \rangle_{0,\beta}^{Y,D^+}$, which vanishes by Lemma 2.8 since $\beta_2 \cdot D' = \beta \cdot D = 0$.

Therefore, suppose that there is a nonzero term with nontrivial Γ_1 -curve and with the Γ_2 -curve in the nonzero class β_2 . As in Lemma 2.8 denote by $\beta_{2,j} := \ell_j f + \alpha'_j$ the classes of its connected components, where $\alpha'_j \in H_2(D')$. If $\alpha'_j \neq 0$ then $\ell_j \neq 0$ because the whole curve is connected. Hence Lemma 2.8 implies that $\beta_{2,j} \cdot D' = \ell_j + \alpha'_j \cdot D' < 0$. Therefore $k_j := -\alpha'_j \cdot D' \geq 2$.

We now show that this is ruled out by our dimensional hypothesis. Suppose that the $\beta_{2,j}$ -curve has r_j relative constraints with total (complex) dimension δ_{bj} . Then, because $c_1^Y(\alpha'_j) = c_1^X(\alpha'_j)$, the dimensional condition

$$n + 2\ell_j + c_1^X(\alpha'_j) + r_j - 3 + \delta_{bj} - (\ell_j - r_j) - r_j n = 0$$

must hold. Suppose now that $n \geq 3$. Since $k_j > \ell_j \geq r_j \geq 1$ the LHS decreases strictly if we replace ℓ_j by r_j , δ_{aj} by 0 and $c_1^X(\alpha'_j)$ by $(n-3)(k_j-2)$. Hence (dropping the subscripts j) we must have

$$n + 3r - 3 + (n-3)(k-2) - rn = (n-3)(k-r-1) < 0.$$

which is impossible. A similar argument works when $n = 2$. This proves (i).

Now consider (ii). There is a term in the decomposition formula for $\langle a_1, \dots, a_k \rangle_{0,\beta}^X$ in which a connected Γ_1 -curve in class β meeting D transversally in $d = \beta \cdot D$ arbitrary points is joined to d disjoint fibers in Y . Again we must show that there are no other nonzero terms. It follows from Lemma 2.8 (ii) that any component in Γ_2 consisting only of a fiber class $d_i f$ must have $d_i = 1$. Hence, if there is another term one of the components of the Γ_2 curve must lie in some class $\beta_{2,j} = d_j f + \alpha'_j$ where $\alpha'_j \neq 0$. But these can be ruled out just as in case (i). $\square$

Remark 3.3. (i) As noted by Hu [5], this lemma applies when D is the blow up of a point or of a curve S with $c_1(S) \geq 0$.

(ii) Maulik–Pandharipande [14] show that the invariant $\langle |b_1^*, \dots, b_r^* \rangle_{0,\ell f + \alpha', \underline{d}}^{Y,D^+}$ may be calculated in terms of suitable descendent invariants of D' in class α' . Hence in the above lemma it also suffices to restrict the assumption on $\alpha \cdot D$ to classes α for which the descendent GW invariants in D do not all vanish.

We now give some examples to illustrate the difference between the absolute and relative invariants.

It is easy to find examples with $\beta \cdot D < 0$ for which the two invariants are different since the relative invariant vanishes by definition, while the absolute invariant might not. For example, consider $X = S^2 \times S^2$ with $\alpha_1 := [S^2 \times pt]$ and $\alpha_2 := [pt \times S^2]$, and let D be the antidiagonal in class $\alpha_1 - \alpha_2$. Then

$$\langle pt | \rangle_{0,\alpha_1}^{X,D} = 0, \quad \langle pt \rangle_{0,\alpha_1}^X = 1.$$

A similar phenomenon may happen when $\beta \cdot D = 0$. We first give a higher genus case and then in Proposition 3.7 a genus zero example.

Example 3.4. *Different absolute and relative genus one invariants when $\beta \cdot D = 0$:*

Let $X = \mathbb{P}(L \oplus \mathbb{C})$ where $L \rightarrow T^2$ is a holomorphic line bundle of degree 1 and put $D = \mathbb{P}(L \oplus \{0\})$ the section of self-intersection -1 . Let $\beta = D + f$, the class of the section $D^+ := \mathbb{P}(\{0\} \oplus \mathbb{C})$. Then $\beta \cdot D = 0$. The dimension of the moduli space $\overline{\mathcal{M}}_{1,1}(X, \beta, J)$ of holomorphic tori in X with one marked point is 2, and so there should be a finite number of such tori through a generic point. We claim that:

$$(3.2) \quad \langle pt \rangle_{1,\beta}^X = 2; \quad \langle pt | \rangle_{1,\beta}^{X,D} = 1.$$

If J is the obvious complex structure on the ruled surface X then there is a 1-parameter family of tori $z \mapsto [\lambda\sigma(z) : 1]$ in class β , where σ is a holomorphic section of L and $\lambda \in \mathbb{C}$. All these tori go through the point p where D^+ meets the fiber over the point z_0 where σ vanishes. The Riemann–Roch theorem implies that these are all regular curves because they are embedded with self-intersection > 0 . There are infinitely many such tori through p , but just one through any other point $q \in X \setminus D$. However there is also a reducible β -torus through each point, the union of D with a fiber of $X \rightarrow D$. Thus this standard J is *not* regular for tori in class β . In fact, one can show that $\langle pt \rangle_{1,\beta}^X = 2$, i.e. the family of reducible curves also contributes 1 to this invariant. (This can be checked by comparing with the Seiberg–Witten invariants; cf Li–Liu [12]. Or one could make a gluing argument as in Example 2.9.)

To calculate the relative invariant $\langle pt | \rangle_{1,\beta}^{X,D}$, consider the relative moduli space $\overline{\mathcal{M}}_{1,\beta}(X, D; J)$ for the standard complex structure on X . (Note that we must use an almost structure for which D is holomorphic.) Then the prolongation $Q = \mathbb{P}(\mathbb{C} \oplus L^*)$ can be identified with the complex manifold X in such a way that (D_0, D_∞) correspond to (D^+, D) . Hence the only holomorphic sections of Q are tori through the point z_D in D_0 lying over z_0 . Therefore there is only *one* curve in $\overline{\mathcal{M}}_{1,\beta}(X, J)$ with two levels; it consists of the fiber over z_0 together with a torus in Q through z_D . Therefore, there is just one element of $\overline{\mathcal{M}}_{1,\beta}(X, D; J)$ that goes through a generic point of $X \setminus D$, and this element is regular. Hence $\langle pt | \rangle_{1,\beta}^{X,D} = 1$. This proves (3.2).

Similarly, $\langle |pt \rangle_{1,\beta,(1)}^{X,D^+} = 1$. The best way to see this is to take a generic complex structure for which D^+ is holomorphic. Then because the absolute GW invariant is 2 there is one more holomorphic torus through each point of D^+ . This set of tori form the main stratum of the moduli space. The curve D^+ itself does not appear and, as before, there is only one section in Q . Since this may well not go through the given point constraint on $D_{1\infty}$ it does not contribute to the invariant.

On the other hand $\langle pt | D^+ \rangle_{1,\beta,(1)}^{X,D^+} = 2$, since if J is a generic complex structure for which D^+ is holomorphic there are precisely two holomorphic tori in class β through any point not on D^+ that each meet D^+ transversally.

We now show that *these results are consistent with the decomposition formula*. Let us calculate $\langle pt \rangle_{1,\beta}^X$ by thinking of X as the connected sum of (X, D) with the ruled surface (Y, D^+) , defined as in Proposition 3.2. Since (Y, D^+) can be identified with (X, D^+) we have $\langle |pt \rangle_{1,\beta,(1)}^{Y,D^+} = 1$. Therefore, if we put the point constraint into X , besides the term

$$\langle pt | \rangle_{1,\beta}^{X,D} = 1$$

with $\Gamma_2 = \emptyset$, there also is a term in the decomposition formula in which the curve in (X, D) is the sphere fiber through pt , and the curve in the ruled surface Y is a torus in class β meeting D^+ in a point constraint. Thus this contribution is the product

$$\langle pt | D \rangle_{0,f,(1)}^{X,D} \langle |pt \rangle_{1,\beta,(1)}^{Y,D^+} = 1.$$

Note that here the constraints along the divisor $D \equiv D^+$ are dual, as required. Thus

$$\langle pt \rangle_{1,\beta}^X = \langle pt | \rangle_{1,\beta}^{X,D} + \langle pt | D \rangle_{0,f,(1)}^{X,D} \langle |pt \rangle_{1,\beta,(1)}^{Y,D^+} = 2.$$

One can also apply the decomposition formula with the point constraint in Y . Then there is just one nonzero term, and we find

$$\langle pt \rangle_{1,\beta}^X = \langle |pt \rangle_{0,f,(1)}^{X,D} \langle pt | D^+ \rangle_{1,\beta,(1)}^{Y,D^+} = 2,$$

as before. □

3.1. A genus zero example. In this section we give a genus zero example where $\beta \cdot D = 0$ but the relative and absolute invariants are different. Throughout X is the blow up of $\mathbb{P}^4$ at two points, and D is the proper transform of a hyperplane through these two points. Thus $D = \mathbb{P}^3 \# 2\overline{\mathbb{P}}^3$ as in Example 2.9. As a homology class in X , $D = H^X - E_1^X - E_2^X$ where H^X is the hyperplane in X and E_j^X are the exceptional divisors. We define λ to be the class of the line in $\mathbb{P}^4$ and ε_j to be the class of the line in E_j , considered where appropriate as elements of $H_2(D)$ or $H_2(X)$. We shall use the self-dual basis

$$(3.3) \quad pt, \lambda, \varepsilon_1, \varepsilon_2, \lambda^* := \pi, \varepsilon_1^*, \varepsilon_2^*, pt^* = D$$

for $H_*(D)$. Thus $\pi := D \cap H^X$ and $\varepsilon_j^* = -D \cap E_j^X$ are disjoint (and equal to the classes $H, -E_j$ in the notation of Example 2.9).

We need two auxiliary lemmas. The first is due to Gathmann. It compares certain Gromov–Witten invariants in X and $\mathbb{P}^4$, and expresses the fact that these invariants are enumerative, i.e. they count what one expects them to count.

Lemma 3.5. *Let $X = \mathbb{P}^4 \# 2\overline{\mathbb{P}}^4$ be as above and $a_1, \dots, a_m \in H_{<8}(\mathbb{P}^4) \subset H_{<8}(X)$.*

(i) *For any $k > 0$, $\langle a_1, \dots, a_m \rangle_{0,k\lambda}^X = \langle a_1, \dots, a_m \rangle_{0,k\lambda}^{\mathbb{P}^4}$.*

(ii) *For any $k > 0$ and $i = 1, 2$, $\langle a_1, \dots, a_m \rangle_{0,k\lambda-\varepsilon_i}^X = \langle pt, a_1, \dots, a_m \rangle_{0,k\lambda}^{\mathbb{P}^4}$.*

(iii) *For any $k > 0$, $\langle a_1, \dots, a_m \rangle_{0,k\lambda-\varepsilon_1-\varepsilon_2}^X = \langle pt, pt, a_1, \dots, a_m \rangle_{0,k\lambda}^{\mathbb{P}^4}$.*

The first part of the next lemma is also well known; we give the proof for completeness.

Lemma 3.6. *Let X, D be as above. Then*

- (i) $\langle pt, pt, \lambda, \pi, \pi, \pi \rangle_{0,2\lambda}^{\mathbb{P}^4} = 4.$
- (ii) $\langle pt, pt, \pi, \pi, \pi \mid \lambda, D \rangle_{0,2\lambda,(1,1)}^{X,D} = 8.$

Proof. (i) Put the constraints $pt, pt, line$ in general position and let P be the 3 space containing them. Then every conic (i.e. degree 2 holomorphic curve) through these constraints intersects P in three points and so must lie entirely in P . It follows readily that

$$\langle pt, pt, \lambda, \pi, \pi, \pi \rangle_{0,2\lambda}^{\mathbb{P}^4} = \langle pt, pt, \lambda, \lambda, \lambda, \lambda \rangle_{0,2\lambda}^{\mathbb{P}^3}.$$

Now calculate the invariant by applying the splitting rule (cf. [20, Ch 7.5]) to

$$I = \langle pt, pt, \pi, \pi; \lambda, \lambda, \lambda \rangle_{0,2\lambda}^{\mathbb{P}^3, \{1,2,3,4\}},$$

where the superscript $\{1,2,3,4\}$ means that the cross ratio of the first four marked points is fixed and we have dropped the subscript H . If the constraints are split as pt, pt and π, π then we find

$$\begin{aligned} I &= \langle pt, pt, \lambda, \lambda, \lambda, \lambda \rangle_{0,2\lambda}^{\mathbb{P}^3} \langle \lambda^*, \pi, \pi \rangle_{0,0}^{\mathbb{P}^3} + \langle pt, pt, \pi \rangle_{0,\lambda}^{\mathbb{P}^3} \langle \pi^*, \lambda, \lambda, \lambda \rangle_{0,\lambda}^{\mathbb{P}^3} \\ &= \langle pt, pt, \lambda, \lambda, \lambda, \lambda \rangle_{0,2\lambda}^{\mathbb{P}^3} + 2, \end{aligned}$$

because $\lambda^* = \pi$ and $\langle \lambda, \lambda, \lambda, \lambda \rangle_{0,\lambda}^{\mathbb{P}^3} = 2$ by [20, Example 7.1.16]. On the other hand if they are split with pt, π in each factor then we get

$$I = 6 \langle pt, \pi, \pi, \lambda, \lambda \rangle_{0,\lambda}^{\mathbb{P}^3} \langle pt, \pi, \pi^*, \lambda \rangle_{0,\lambda}^{\mathbb{P}^3} = 6,$$

where the second equality uses the fact that $\langle pt, \lambda, \lambda \rangle_{0,\lambda}^{\mathbb{P}^3} = 1$ by [20, Example 7.1.14]. This proves (i).

Now consider (ii). We first claim that

$$\langle pt, pt, \pi, \pi, \pi \mid \lambda, D \rangle_{2\lambda,(1,1)}^{X,D} = \langle pt, pt, \pi, \pi, \pi, \lambda, D \rangle_{2\lambda}^X.$$

This does not immediately follow from Proposition 3.2(ii) since D does not satisfy the conditions of this lemma for all classes α . However, all we need is that c_1^X is sufficiently positive on the classes α that could appear in a decomposition (β_1, β_2) of the class 2λ . But this condition is satisfied because, if we write $\beta_2 = \ell f + s' \lambda' - m'_1 \varepsilon'_1 - m'_2 \varepsilon'_2$, we must have $m'_1 = m'_2 = 0$. (Otherwise there would be a connected component in Γ_1 in some class $p\varepsilon_j$ for $p \geq 1$. But such a class cannot be controlled by the available absolute constraints since π can be represented by a pseudocycle that is disjoint from

the ε_j -curves.⁶⁾ Therefore

$$\begin{aligned} \langle pt, pt, \pi, \pi, \pi | \lambda, D \rangle_{2\lambda, (1,1)}^{X,D} &= \langle pt, pt, \pi, \pi, \pi, \lambda, D \rangle_{2\lambda}^X \\ &= 2 \langle pt, pt, \pi, \pi, \pi, \lambda \rangle_{2\lambda}^X, \end{aligned}$$

where the last equality holds by the divisor axiom and the fact that $2\lambda \cdot D = 2$. Now apply Lemma 3.5 to reduce this to an invariant in $\mathbb{P}^4$ and use (i). $\square$

We now give an explicit example in genus zero where the absolute and relative invariants differ.

Proposition 3.7. *Let (X, D) and π be as in the previous lemma, and let $\beta = 4\lambda - 2\varepsilon_1 - 2\varepsilon_2$. Then*

$$\langle pt, pt, \pi, \pi, \pi \rangle_{\beta}^X \neq \langle pt, pt, \pi, \pi, \pi | \rangle_{\beta}^{X,D}.$$

Proof. The difference between these invariants is

$$\sum_{\beta_2 \neq 0, b_i, a_j, \underline{d}} \langle pt, pt, a_1, \dots, a_q | b_1, \dots, b_r \rangle_{\beta_1, \underline{d}}^{\Gamma_1, X, D} \langle a'_1, \dots, a'_{q'} | b_1^*, \dots, b_r^* \rangle_{\beta_2, \underline{d}}^{\Gamma_2, Y, D^+}$$

where we sum over all possible relative constraints and all placings of the absolute constraints π, π, π (which we decompose as in Lemma 2.8). Thus $q + q' = 3$.

There is one nonzero term in this sum with $\beta_1 = 2\lambda$ and $\beta_2 = 2\alpha' + 2f$. Here we take the Γ_1 curve to be a connected 2λ curve through all the absolute constraints and the relative constraints $b_1 := D, b_2 := \lambda$, and the Γ_2 curve to be disconnected, with one component equal to a fiber through a point in $b_1^* = pt$ and the other a sphere in class $2\alpha' + f$ and through the relative constraint $b_2^* =: \pi$. The Γ_2 factor equals $1/4$ by Example 2.9. Further, by Lemma 3.6

$$\langle pt, pt, \pi, \pi, \pi | \lambda, D \rangle_{0, 2\lambda, (1,1)}^{X,D} = 8.$$

Therefore this term contributes 2 to the sum.

Since curves in class 2α do not meet $D \cap E_j$,

$$\langle pt, pt, \pi, \pi, \pi | \varepsilon_j, D \rangle_{0, 2\lambda, (1,1)}^{X,D} = 0, \quad j = 1, 2,$$

where we have used the notation of equation (3.3). Moreover, although

$$\langle pt, pt, \pi, \pi | pt, D \rangle_{0, 2\lambda, (1,1)}^{X,D} = \langle pt, pt, \pi, \pi, pt \rangle_{0, 2\lambda}^X = 1,$$

the corresponding Γ_2 invariant vanishes. For, this would have to consist of a fiber together with a connected curve in class $f + 2\alpha'$. This is nonzero only if there is a $\mathbb{C}^*$ invariant representative of $f + 2\alpha'$ through the remaining absolute constraint. But we can arrange that there are no such curves: by Remark 3.1 (iii) the constraint has the form $\pi^{-1}(\lambda')$, where λ' is a line in D' that can certainly be chosen disjoint from the α' curve. A similar argument rules out the case when Γ_1 consists of two components each

⁶ One must be careful here since there are nonzero invariants in class ε_j . For example, if $\varepsilon_j, \varepsilon_j^*$ denote the line and 2-plane in the exceptional divisor E_j^X in X $\langle \varepsilon_j^*, \varepsilon_j^* | \varepsilon_j \rangle_{0, \varepsilon_j}^{X,D} = \langle \varepsilon_j, \varepsilon_j, pt \rangle_{0, \varepsilon_j}^{E_j^X} = 1$.

in class λ , since such Γ_1 cannot go through all the absolute constraints.⁷ Therefore there are no other contributions with Γ_1 a connected curve in class 2λ .

We now show that there are no other nonzero terms in the sum. To see this, consider the possible choices for β_2 . By Lemma 2.8, each connected component of a nonzero Γ_2 factor must either lie in class f or in some class $\beta_{2,i}$ such that $\beta_{2,i} \cdot D' < 0$. Since $\beta \cdot D < 0$ there must be at least one term of the latter form.

Consider such a term and write $\beta_{2,i} = k_i f + s'_i \lambda' - m'_{i1} \varepsilon'_1 - m'_{i2} \varepsilon'_2$. Then since $k_i \geq 1$ we need $s'_i - m'_{i1} - m'_{i2} \leq -2$. Since the class $s'_i \lambda' - m'_{i1} \varepsilon'_1 - m'_{i2} \varepsilon'_2$ must be representable in D we need either $s'_i > 0$ or $s'_i = 0$ and $m'_{i1} \leq 0$. But the latter possibility makes $s'_i - m'_{i1} - m'_{i2} \geq 0$. Therefore $s'_i > 0$.

We next claim:

if the class $s\lambda - m_1\varepsilon_1 - m_2\varepsilon_2$ with $s > 0$ is represented by a holomorphic sphere S in D then either $m_1 = m_2 = s$ or $0 \leq m_j$ and $m_1 + m_2 \leq s$.

Proof of Claim: In the notation of Example 2.9,

$$(s\lambda - m_1\varepsilon_1 - m_2\varepsilon_2) \cdot_D (H - E_1 - E_2) = s - m_1 - m_2.$$

If this is negative then S lies entirely in the plane $H - E_1 - E_2 \cong \mathbb{P}^2 \# 2\overline{\mathbb{P}}^2$. Hence it must either be a multiple of the unique curve in class $\alpha = \lambda - \varepsilon_1 - \varepsilon_2$ or have nonnegative intersection with it. The claim follows.

The connected component $\beta_{2,i}$ might be a union of spheres, but, by the claim, the homology class of each such sphere is $s_i\alpha$, $m\varepsilon_j$ or $s\lambda - m_1\varepsilon_1 - m_2\varepsilon_2$ with $s \geq m_1 + m_2 \geq 0$. Since the classes $s_i\alpha$ are the only ones with negative intersection with D' there must be at least one of these. Moreover, we need $s_i \geq 2$ for the intersection to be negative and also $\sum_i s_i \leq 3$ since the Γ_1 component goes through the two point constraints.

If $\sum_i s_i = 3$, then the Γ_1 component must be

$$\langle pt, pt \mid D \rangle_{0,\lambda}^{X,D}.$$

Therefore the Γ_2 component is connected and must go through the absolute constraints. As noted above, these have the form $\pi^{-1}(\lambda')$ where we may take $\lambda' \subset D'$ to be disjoint from the exceptional divisors E'_1, E'_2 . Further $\beta_2 = f + 3\alpha + \varepsilon_1 + \varepsilon_2$. Therefore its only holomorphic and $\mathbb{C}^*$ invariant representatives consist of the union of a fiber with a triply covered α -curve and some curves in $E'_1 \cup E'_2$. Since these do not meet the absolute constraints, there are no contributions of this form.

Therefore Γ_2 has just one nonfiber component, and this must lie in class $kf + 2\alpha$. Hence $\beta_1 = 2\lambda$. But we have already seen that there is only one term of this form. This completes the proof. $\square$

Remark 3.8. One could calculate $\langle pt, pt, \pi, \pi, \pi \rangle_{\beta}^X$ by Gathmann's algorithm, but the argument is quite complicated and is unnecessary for the present purpose. Note that

⁷ It is possible to use up two absolute constraints because

$$\langle pt, \pi, \pi \mid \pi \rangle_{0,\lambda}^{X,D} = \langle pt, \pi, \pi, \pi \rangle_{0,\lambda}^{\mathbb{P}^4} = \langle pt, \pi, \lambda, \lambda \rangle_{0,\lambda}^{\mathbb{P}^3} = 1.$$

the reason why the two invariants differ in the case considered here is precisely the same as the reason why many of the invariants considered by Gathmann are not enumerative, namely the class β can be represented by stable maps with components in D that are regular in D but not in X .

REFERENCES

- [1] F. Bourgeois, Y. Eliashberg, H. Hofer, K. Wysocki and E. Zehnder, Compactness results in Symplectic Field theory, *Geom. and Top.* **7** (2003), 799–888.
- [2] A. Gathmann, Gromov–Witten invariants of blowups, *J. Algebraic Geometry* **10** (2001), 399–432.
- [3] H. Hofer, A general Fredholm theory and applications, arxiv:math/0509366.
- [4] H. Hofer, C. Wysocki, and E. Zehnder, A General Fredholm Theory I: a splicing-based differential geometry, arxiv:math/0612604.
- [5] J. Hu, Gromov–Witten invariants of blow ups along points and curves, *Math. Z* **233** (2000), 709–39.
- [6] J. Hu, T.-J. Li and Yongbin Ruan, Birational cobordism invariance of uniruled symplectic manifolds, arxiv:math/0611592, to appear in *Invent. Math.*.
- [7] J. Hu and Yongbin Ruan, Positive divisors in symplectic geometry, arxiv:math/0802.0590
- [8] E. Ionel and T. Parker, The Symplectic Sum formula for Gromov–Witten invariants, *Ann. of Math.* **159** (2004), 935–1025.
- [9] Hsin-Hong Lai, Gromov–Witten invariants of blow-ups along submanifolds with convex normal bundles, arxiv:math/0710.3968.
- [10] Y.-P. Lee and R. Pandharipande, A reconstruction theorem in Quantum Cohomology and Quantum k -theory, arxiv:math/0104084.
- [11] An Li and Y. Ruan, Symplectic surgery and Gromov–Witten invariants of Calabi–Yau 3-folds, *Invent. Math.* **145** (2001), 151–218.
- [12] T.J. Li and A. Liu, Symplectic structure on ruled surfaces and generalized adjunction formula, *Math Research Letters* **2** (1995), 453–471.
- [13] T.-J. Li and Y. Ruan, Uniruled symplectic divisors, arxiv:math/0711.4254
- [14] D. Maulik and R. Pandharipande, A topological view of Gromov–Witten theory, to appear in *Topology*, arxiv:math/0412503.
- [15] D. McDuff, Almost complex structures on $S^2 \times S^2$, *Duke Math. Journal* **101** (2000), 135–177.
- [16] D. McDuff, Quantum homology of Fibrations over S^2 , *International Journal of Mathematics*, **11**, (2000), 665–721.
- [17] D. McDuff, Branched manifolds, groupoids, and multisections, SG/0509664, *Journal of Symplectic Geometry*.
- [18] D. McDuff, Hamiltonian S^1 -manifolds are uniruled, arXiv:math/0706.0675, to appear in *Duke Math. Journ.*.
- [19] D. McDuff and S. Tolman, Topological properties of Hamiltonian circle actions, SG/0404338, *International Mathematics Research Papers* vol **2006**, Article ID 72826, 1–77.
- [20] D. McDuff and D.A. Salamon, *J-holomorphic curves and symplectic topology*. Colloquium Publications **52**, American Mathematical Society, Providence, RI, (2004).
- [21] A. Zinger, Enumerative vs. Symplectic invariants and Obstruction bundles, arxiv:math/0201255, *Journ. Symp. Geom.*
- [22] A. Zinger, A comparison theorem for Gromov–Witten invariants in the symplectic category, arxiv:math/0807.0805
- [23] A. Zinger, in preparation.

DEPARTMENT OF MATHEMATICS, BARNARD COLLEGE, COLUMBIA UNIVERSITY, NY 10027, USA
E-mail address: `dusa@cpw.math.columbia.edu`